

Clinical utility of kallikrein-related peptidases (KLK) in urogenital malignancies

Julia Dorn¹; Jane Bayani^{2,3}; George M. Yousef^{2,4}; Feng Yang¹; Viktor Magdolen¹; Marion Kiechle¹; Eleftherios P. Diamandis^{2,3}; Manfred Schmitt¹

¹Clinical Research Unit, Department of Obstetrics and Gynecology, Klinikum rechts der Isar, Technische Universität München, Munich, Germany; ²Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Ontario, Canada; ³Department of Pathology and Laboratory Medicine, Mount Sinai Hospital, Toronto, Ontario, Canada; ⁴Department of Laboratory Medicine and the Keenan Research Centre in the Li KaShing Knowledge Institute, St Michael's Hospital, Toronto, Ontario, Canada

Summary

Kallikrein-related peptidases (KLK), which represent a major tissue-associated proteolytic system, stand for a rich source of biomarkers that may allow molecular classification, early diagnosis and prognosis of human malignancies as well as prediction of response or failure to cancer-directed drugs. International research points to an important role of certain KLKs in female and male urogenital tract malignancies, in addition to cancers of the lung, brain, skin, head and neck, and the gastrointestinal tract. Regarding the female/male urogenital tract, remarkably, all of the KLKs are expressed in the normal prostate, testis, and kidney whereas the uterus, the ovary, and the urinary bladder are expressing a limited number of KLKs only. Most of the information re-

garding KLK expression in tumour-affected organs is available for ovarian cancer; all of the 12 KLKs tested so far were found to be elevated in the malignant state, depicting them as valuable biomarkers to distinguish between the normal and the cancerous phenotype. In contrast, for kidney cancer, a series of KLKs was found to be down-regulated, while other KLKs were not expressed. Evidently, depending on the type of cancer or cancer stage, individual KLKs may show characteristics of a Janus-faced behaviour, by either expanding or inhibiting cancer progression and metastasis.

Keywords

Cancer, proteases, endometrium, ovary, prostate

Correspondence to

Prof. Dr. Manfred Schmitt
Klinische Forschergruppe der Frauenklinik der Technischen Universität München
Ismaninger Strasse 22, 81675 München, Germany
Tel.: +49 89 4140 2449, Fax: +49 89 4140 7410
E-mail: manfred.schmitt@LRZ.tum.de

Received: March 10, 2013

Accepted after major revision: May 25, 2013

Prepublished online: June 13, 2013

doi:10.1160/TH13-03-0206

Thromb Haemost 2013; 110: ■■■■

Introduction

In the United States of America (USA), for the year 2012, 848,170 new cancer cases were predicted for the male population, and 790,740 new cases for women. Of that, 251,900 new cancer cases (29.7%) would affect the male reproductive system compared to 88,750 (11.2%) cases affecting the female reproductive system (1, 2). For the urinary system, an estimated 97,610 new cases (11.5%) were estimated for men and 43,530 new cases (5.5%) for females. All of these cancers can invade nearby organs or metastasise through the bloodstream or lymphatic system to other, distant parts of the body.

Gynaecological cancers include the more common cancers of the corpus uteri (endometrium), cervix uteri, and the ovaries, in addition to the rare cancers of the vagina, vulva, and the fallopian tubes. In the western world, the incidence of gynaecological malignancies is highest for endometrial cancer, followed by cervical and ovarian cancer, while the mortality rate is quite opposite: highest for ovarian cancer, followed by cervical and endometrial cancer (3). Staging of all of the gynaecological malignancies is performed according to the guidelines of the International Federation of Gynaecology and Obstetrics (4). With an 349,510 new cases estimated, cancers of the male urogenital tract encompassing those of

the prostate, testis, penis, kidney, and bladder, accounted for over 41.2% of all new cases estimated. Within that group, prostate cancer is the most frequent cancer in men (estimated 241,740 new cases, 24.5%) and testicular cancer (estimated 8,590 new cases, 1.1%) most common in young men (1, 2).

Current research aims at understanding the biology underlying these disease processes, especially understanding the complex molecular interactions within a tumour and molecular actions of drugs. To identify targets for new and novel therapies remains important to improve our knowledge of the pathobiology of these cancer diseases and associated chemoresistance. Cancer biomarkers such as the kallikrein-related peptidase family members (KLK) may help to identify patients at risk and to see how well a body afflicted with cancer responds to cancer therapy and may even help to decide how aggressive the cancer treatment should be (5, 6).

Expression of KLKs (kallikrein-related peptidases) in the urogenital system

KLK1–15 comprise 15 highly conserved serine proteases with similar structural characteristics and a wide spectrum of func-

tional properties (7). In the female reproductive system, the uterus, endometrium, cervix uteri, and the ovarian and fallopian tube tissues show a variable pattern of expression among the 15 KLK family members (8). During the menstrual cycle, the innermost glandular layer of the uterus, the endometrium, grows to a thick glandular tissue layer, and shows expression of KLKs for most of the KLKs, both in the proliferative and the secretory phase. Variable expression of KLKs is observed in the mucin-secreting epithelium of the endocervix and the tubular cervical glands as well (9, 10). Different from these tissues, a few KLKs are highly expressed within the ovary and/or the fallopian tube, others are generally absent or only weakly expressed (11–14) (► Tables 1–3).

In the male reproductive system, prostatic tissues show high expression of selected KLKs (8, 15) (► Table 4 and ► Table 5). Assessment of the topographical distribution of several of the KLK antigens by immunohistochemistry showed mostly cytoplasmic staining in the luminal secretory cells of the prostatic epithelium, in basal cells, and in foci of basal cell hyperplasia. As an exception, KLK4 appears to be located to the nucleus as well (16, 17). The epithelium of the rete testis, ejaculatory ducts, seminal vesicles, ductus deferens, epididymis efferent ductules, penile urethra and Littre's and Cowper's glands express KLKs as well; the spermatogenic epithelium in the testis shows variable expression of KLKs with spermatogonia being the predominant KLK expressing cell population (11–14, 17).

The urinary system includes the kidneys, bladder, and ureter. In the normal kidney, there is relatively low mRNA expression of several of the KLKs but high expression of others (► Table 6) (8, 12, 13). In the normal urinary bladder, so far, expression patterns of only a few of the KLKs have been investigated (8). Surprisingly,

urine contains only few KLKs, e.g. KLK1, 3, 4, 9, and 12, present at low levels (8).

Deregulated expression, secretion, and function of various KLK family members have been reported for solid malignant tumours of the urogenital tract, including kidney cancer, cervical cancer and hormone-dependent cancers of the endometrium, ovary, prostate, and the testis. In these cancers, KLKs may serve as diagnostic, prognostic, or therapy-response cancer biomarkers and may even help to distinguish between various types of cancer when morphology is similar (18). Still, there is not a uniform pattern concerning mRNA or protein expression of the various KLKs under malignant conditions. KLKs are downregulated in kidney cancer but upregulated in ovarian cancer, compared to their non-malignant cellular counterparts (15, 17, 18); while some of the KLKs are downregulated in prostate cancers, others are upregulated (15, 17, 19).

Clinical utility of KLKs in cancers of the female reproductive tract

KLKs in endometrial cancer

Endometrial cancer originates from the inner glandular layer of the uterus. The leading and early symptom of endometrial cancer is vaginal discharge or abnormal bleeding. Because of this, endometrial cancer is often diagnosed in an early stage, giving hope for a favourable clinical course of the disease. The metabolic syndrome, comprising obesity, diabetes, and dislipidaemia, as well as infertility, polycystic ovarian syndrome, Lynch syndrome, and early menarche and late menopause are known risk factors for en-

	Normal (uterus)		Normal (endometrium)		Cancerous+		References
	mRNA	Protein	mRNA	Protein	mRNA	Protein	
KLK1	Low	Low	Expressed	Expressed	↓	ND	8, 44, 142, 143
KLK2	Absent	Absent	Expressed	ND	ND	ND	8, 144
KLK3	Low	Absent	Expressed	Expressed	ND	ND	8, 144, 146
KLK4	Absent	Low	ND	Expressed	ND	↑	8, 25, 44
KLK5	Absent	Absent	ND	Expressed	ND	ND	8, 9, 16, 27
KLK6	Absent	Low	Expressed	Expressed	↑	ND	8, 9, 12, 16, 26, 27, 44, 145
KLK7	Absent	Absent	ND	Expressed	ND	ND	8, 16, 27, 44
KLK8	Absent	Absent	Expressed	Expressed	↑	↑	8, 10
KLK9	Absent	Moderate	ND	ND	ND	ND	6, 8
KLK10	Moderate	Absent	Expressed	Expressed	↑	ND	8, 13, 16, 44, 145
KLK11	Low	Low	ND	Expressed	ND	ND	8, 9, 16, 27, 44
KLK12	Low	Moderate	ND	Expressed	ND	ND	8, 9, 16, 27, 44
KLK13	Absent	Low	ND	Expressed	ND	ND	8, 9, 16, 27
KLK14	Low	Low	ND	Expressed	ND	ND	8, 16, 56
KLK15	Absent	Absent	ND	ND	ND	ND	8

↑ = Increased, ↓ = decreased compared to normal tissue. ND = Not determined, +Endometrial cancer.

Table 1: KLKs present in normal and tumour tissues of patients afflicted with endometrial cancer.

ometrial cancer (20, 21). Therapy of choice of endometrial cancer is hysterectomy with bilateral salpingo-oophorectomy, often associated with pelvic and paraaortal lymphadenectomy and/or followed by adjuvant radiotherapy. Chemotherapy or endocrine therapy is primarily administered in advanced stages of endometrial cancer (22). Currently, no effective serological or tissue biomarkers exist to classify patients at risk. The serum biomarker CA125 (a tumour cell surface mucin encoded by the *MUC16* gene) can be used in single cases to monitor effect of therapy, but one has to keep in mind that only 25% of the endometrium cancer patients are positive for CA125 (23).

Published data are available regarding KLK mRNA and/or protein expression in tissue extracts of normal uterus/endometrium and its malignant counterparts (8) (► Table 1). Immunoenzymometric testing of tissue extracts of the uterus revealed that eight KLK proteins were expressed (KLK1, 4, 6, 9, and 11-14), seven were not. At the mRNA level, six of the KLKs were detected (KLK1, 3, 10-12, and 14). Limited data are available for KLK expression in the normal endometrium, at the mRNA level and at the protein level, as assessed by immunohistochemistry. Six KLK mRNAs (KLK1-3, 6, 8, and 10) were found to be expressed, for the other nine KLKs no mRNA expression data have been reported. Assessment by immunohistochemical staining demonstrated protein expression of 12 KLKs (KLK1, 3-8, 10-14), no data are available regarding protein expression in the normal endometrium of the other three KLKs (16). Not much of published information is available regarding the mRNA/protein expression patterns of KLKs in endometrial carcinomas. At the mRNA level, KLK1 was found to be downregulated, KLK6, 8, and 10 are upregulated. KLK4 and 8 proteins are upregulated; no data were presented in the scientific literature regarding protein expression of the other 13 KLKs.

KLK4, which is overexpressed in endometrial cancer compared to normal and hyperplastic endometrium as assessed by immunohistochemistry, is upregulated by the steroid hormones oestrogen and progesterone (24, 25). In contrast, KLK6 has emerged as a marker describing a more aggressive type of this cancer disease: in type-II serous papillary endometrial cancer, KLK6 gene expression is significantly elevated compared to normal, benign, and type-I endometrioid carcinoma; KLK6 and 10 proteins are significantly elevated in the blood of patients presenting with serous-papillary endometrial carcinoma (26, 27).

KLKs in cervical cancer

Malignant tumours of the cervix uteri are one of the leading causes of death of young women worldwide, though due to screening programs and successful therapy of pre-malignant lesions and early stages of the disease, cervical cancer has become a rare disease in the industrialised world. Cervical cancer is known to develop stepwise from infection with the human papilloma virus (HPV) and subsequent inefficient immune response to eliminate the virus, followed by cervical dysplasia (CIN I-III), eventually leading into an invasive type of cervical carcinoma (28). One of the most important factors to predict the clinical outcome of the disease is

Table 2: KLKs present in normal and tumour tissues of patients afflicted with cervical cancer.

	Normal		Cancerous		References
	mRNA	Protein	mRNA	Protein	
KLK1	Medium	Low	ND	ND	8
KLK2	Low	Absent	ND	ND	8
KLK3	Low	Absent	ND	ND	8
KLK4	High	Low	ND	ND	8
KLK5	High	Low	ND	ND	8, 9, 16
KLK6	High	Low	ND	ND	8, 9, 16
KLK7	High	Low	ND	↑	8, 16, 31, 32
KLK8	High	Low	ND	↑	8, 34
KLK9	High	Medium	ND	ND	8
KLK10	High	Low	ND	ND	8, 16
KLK11	High	Medium	ND	ND	8, 9, 16
KLK12	Low	Medium	ND	ND	8, 9, 16
KLK13	High	Low	ND	ND	8, 9, 16
KLK14	Medium	Low	ND	ND	8, 16
KLK15	Absent	Absent	ND	ND	8

ND = Not determined, ↑ = Increased, ↓ = Decreased compared to normal tissue.

clinical stage at the time of diagnosis. Screening by regular gynaecological examination, together with the result of the PAP smear test (the microscopical examination of cervical cells stained according to Papanicolaou) can effectively diagnose premalignant lesions or early cancer stages and thus reduce incidence and mortality of cervical cancer. Management of cervical cancer is stage-dependent: the early invasive cervical cancer is a domain of the surgeon, performing a total radical hysterectomy including dissection of the parametries and pelvic lymph nodes, and closure of the vaginal cuff. In the advanced stages of cervical cancer, primary radio-chemotherapy is indicated (29).

To date, tumour markers play a minor role in the management of cervical cancer. For the squamous cell carcinoma type, efficacy of therapy can be monitored by determination of blood level squamous cell carcinoma antigen, or, for the adenocarcinoma-type, the carcinoembryonic antigen (CEA) and CA125. Admittedly, however, for any stage of cervical cancer, no effective prognostic or predictive cancer biomarkers have been established yet (30).

Although it is known that all of the KLKs are expressed in the normal cervix, except KLK15, either at the mRNA or the protein level (8, 16) (► Table 2), no mRNA KLK expression data are available and only for KLK7 and 8 protein data are published with regard to their expression status in the malignant state. For instance, KLK7 is highly expressed in tumour tissue (31) and KLK7 content increases with the severity of cervical lesions, i.e. cervicitis, low-grade cervical intraepithelial neoplasia, high-grade cervical intraepithelial neoplasia, squamous cervical carcinomas, and even cervi-

cal adenocarcinomas (32). Thus, KLK7 could be a useful marker additional to the PAP smear for screening of cervical precursor lesions (32). Interestingly, the endogenous antileukoprotease inhibitor (ALP/SLPI) of KLK7 is significantly reduced in cervical adenocarcinomas when compared with normal endocervical glands, pointing to a regulatory role of KLK7 and ALP/SLPI in the endometrium (33). As shown for a few tissue specimens only, the KLK8 protein is also expressed in cervical carcinomas although these data have not been validated yet (34).

KLKs in ovarian cancer

The general term "ovarian cancer" encompasses those tumours of the ovary originating from the epithelial surface of this organ and accounts for more than 80% of all solid malignant ovarian tumours. Ovarian carcinomas are categorised according to their histology or molecular subtype. The most common histological epithelial subtype is that of serous cystadenocarcinoma, followed by endometrioid and mucinous carcinoma. Epithelial ovarian cancers are classified molecularly into type I and type II (35), with type I ovarian cancers including low-grade serous-papillary, endometrioid, and borderline tumours of low malignant potential. Type II ovarian cancers are more frequent (75%) and encompass high-grade serous carcinomas, undifferentiated carcinomas, and carcinosarcomas.

Prognosis of patients afflicted with ovarian cancer is poor owing to late diagnosis and often inefficient primary debulking surgery, but also because of rapidly developing chemoresistance. Potentially, early detection of ovarian cancer could decrease mortality, but unfortunately, there is no effective screening procedure: neither vaginal ultrasonography nor analysis of serum cancer biomarker CA125 nor any other biomarker analyses at the protein or gene level are specific and sensitive enough for that objective. Thus, ovarian cancer diagnosis is performed by histological inspection of the biopsy. As a result, a number of patients will have laparotomy though only suffering from a previously unknown benign ovarian tumour operable via laparoscopy. In the case of an adnexal mass of unknown dignity, presurgical assessment of CA125 in serum as well as the cancer biomarker tests OVA1 or ROMA can be applied to support a clinician's decision to perform a biopsy or choose the best surgical approach (37).

Yet, about 20% of ovarian cancer patients do not express the CA125 antigen. Similarly, with many false-positive cases for OVA1 and ROMA, it appears to be useful only for preventing a malignant mass to be falsely classified as benign (36). On the other hand, serum biomarker CA125 is established as a useful cancer biomarker to monitor response of CA125-positive ovarian cancer patients to cancer therapy.

For ovarian cancer, the major traditional prognostic factor is FIGO stage at time of diagnosis. For instance, 5-year-survival rate of early FIGO stage I patients is >90% while survival of patients with late FIGO stages III or IV is <25%. Another important traditional prognostic factor is size of residual tumour mass after cytoreductive surgery (38). Regrettably, owing to the lack of suitable biomarkers, at present, ovarian cancer management is not con-

sidering any prognostic or therapy response predicting factors, concerning the course of the disease or a patient's risk to develop disease recurrences (36).

Since KLKs are supposed to contribute to ovarian cancer progression and metastasis, recently, broad scientific interest has focused on these novel cancer biomarkers aiming at exploring their clinical validity (15, 19, 39). For several of the KLKs, both mRNA and protein levels have been reported to be increased in ovarian cancer tissue, compared to levels found in physiological situations (► Table 3).

By enzymometric determination of bulk-extracted normal ovarian tissues, seven KLKs were found to be expressed in the normal ovary (KLK1, 6-8, 10, 11, 14) (8); immunohistochemical staining revealed positivity for KLK5, 12, and 13 as well (16). At the mRNA level, nine KLKs are expressed (KLK1, 6-11, 13, and 14) (8). Ten KLK mRNAs (KLK3-8, 10, 11, 13, 15) were found to be elevated in ovarian cancer tissue compared to normal counterparts, KLK14 is downregulated (40-52). When assessing KLKs at the protein level by immunohistochemistry or ELISA, 12 KLKs (KLK3-11, 13-15) were found to be upregulated in ovarian cancer compared to normal ovarian tissues or other non-malignant gynaecological diseases (38, 44, 53-55).

Five KLK family members (KLK4-7, and 15) are linked to poor prognosis of ovarian cancer patients, two (KLK9 and 14) are associated with a favourable course of cancer disease. For some (KLK8, 10, 11, and 13) the clinical relevance is not clear yet, since, depending on the way of detection (mRNA or protein) and FIGO stage (early or advanced), these KLKs may be a factor to either predict a poor or a good course of cancer disease (56). The prognostic or predictive value of KLK1, 2, and 12 has not been determined yet; KLK3 expression was found not to be correlated with clinical outcome, yet, presence of the *KLK3* rs11084033 allele correlated with poor survival as determined by the Sequenom iPLEX mass array technology (57).

More specifically, *KLK4* mRNA expression in ovarian tumour tissue is associated with untimely disease recurrence and death (58). This feature parallels that of *KLK5* expression in ovarian tumour tissue where elevated *KLK5* mRNA and *KLK5* protein is correlated with shorter disease-free and overall survival (59, 60). Analysis of *KLK6* protein in tumour cytosols using ELISA indicates that *KLK6*-positive tumours are more likely associated with advanced disease, serous histology, and suboptimal debulking. *KLK6* impacts on overall and progression-free survival, especially in ovarian cancer subgroups with a good prognosis at first sight, e.g. with low-grade tumours and optimally debulked patients (61). Further, stromal-cell associated overexpression of *KLK6* protein as assessed by immunohistochemistry is associated with shorter overall and progression-free survival as well (62). High concentrations of *KLK7* mRNA and *KLK7* protein in ovarian tumour tissue were reported to be associated with advanced stage and larger residual tumour mass, or lower nuclear grade disease and those who have been optimally debulked, as well as shorter overall and progression-free survival (53, 63). Increase in *KLK15* mRNA expression in ovarian tumour tissue is also associated with shorter survival (52). Different from *KLK4-7* and *15* expression, *KLK9* and

Table 3: KLKs present in normal and tumour tissues of patients afflicted with ovarian cancer.

	Normal		Cancerous		References
	mRNA	Protein	mRNA	Protein	
KLK1	Moderate	Moderate	ND	ND	8, 130
KLK2	Absent	Absent	ND	ND	8, 130
KLK3	Absent	Absent	↑	↑	8, 50, 57, 130, 147, 148
KLK4	Absent	Absent	↑	↑	8, 43, 58, 84, 130, 149
KLK5	Absent	Present	↑	↑	8, 16, 44, 48, 55, 59, 60, 130, 149–151
KLK6	High	Moderate	↑	↑	8, 16, 42, 44–46, 50, 55, 61, 62, 69, 75, 130, 149, 150, 152–159
KLK7	High	Moderate	↑	↑	8, 16, 42, 44, 45, 47, 48, 53, 63, 85, 130, 149, 150, 160
KLK8	High	Low	↑	↑	8, 42, 44, 45, 50, 65–69, 130, 149, 150
KLK9	Low	Absent	ND	↑	8, 64, 130
KLK10	High	Moderate	↑	↑	8, 16, 38, 42, 44, 45, 49, 50, 55, 70, 86, 130, 149, 150, 155, 158
KLK11	Moderate	Moderate	↑	↑	8, 16, 44, 45, 55, 71, 72, 74, 130
KLK12	Absent	Present	ND	ND	8, 16, 130
KLK13	Low	Present	↑	↑	8, 16, 38, 55, 73, 75, 130
KLK14	Moderate	Low	↓	↑	8, 11, 16, 44, 51, 123, 130, 161
KLK15	Absent	Absent	↑	↑	8, 52, 130, 162

↑ = Increased. ↓ = decreased compared to normal tissue. ND = Not determined.

14 mRNA are considered factors indicating favourable clinical outcome of ovarian cancer patients (51, 64).

Conflicting results exist concerning the clinical utility of four of the KLKs (KLK8, 10, 11, and 13). For instance, expression of KLK8 protein (65, 66) and KLK8 mRNA (67, 68) were determined to be elevated in tumour cytosols of ovarian cancer patients, and expression correlated to a favourable outcome. Contrary to these results, the immunohistochemical AQUA-study by Kountourakis classified KLK8 protein as a marker of poor clinical outcome although results of this image analysis-based technology deserve validation (69). For KLK10, Luo et al. reported that elevated KLK10 protein in ovarian cancer tumour tissue is associated with advanced stage, suboptimal debulking, larger residual tumour mass, serous histology, and shorter progression-free and overall survival (70). These data could not be confirmed by Dorn et al. who reported that low KLK10 protein expression is associated with poor survival (38). In another study, KLK11 and 13 protein levels expressed in ovarian cancer tissues were found to be correlated with early stage disease and favourable overall survival (71–73). This is different from the findings of Shigemasa et al. and White et al. (74, 75) who reported that increased *KLK11* or *13* mRNA expression values are associated with high-grade tumours and poor patient outcome. Evidently, the prognostic value of *KLK11* and *13* mRNA expression differs from the KLK11 and 13 protein expression values.

Since residual tumour mass after primary debulking is one of the most important issues in ovarian cancer, a score to predict surgical outcome based on the clinical factors tumour grading and as-

cites volume plus KLK6 and 13 was established by Dorn et al. (38). We argued that such a score could help to identify those ovarian cancer patients who could be spared the burden of surgery.

Seven KLKs (KLK5–8, 10, 11, and 14) were found to be released into the blood (serum) of early or advanced ovarian cancer patients, and are elevated in comparison to serum levels of healthy individuals, patients with benign diseases of the ovary, or cancer of other origin (11, 59, 66, 76–83). Nine of the KLKs are also enriched in peritoneal ascites of ovarian cancer patients (KLK5–13, not KLK4 or 14) (8, 11, 81, 84). Elevation of KLK5, 6, 10, and 11 proteins in serum are considered markers of poor prognosis (overall or progression-free survival) for the ovarian cancer patient (75, 78–83) while an increase in KLK8 in ascitic fluid indicates a favourable prognosis (66).

When evaluating the potential clinical application of KLK family members in conjunction with the monitoring cancer biomarker CA125, a direct correlation of CA125 and KLK8 expression released into the blood or the ascitic fluid was uncovered (66). Furthermore, El Sherbini et al. (82) presented evidence that the combination of CA125 with KLK6 and 10 renders increased specificity to predict for ovarian cancer, when compared to the clinical impact of each of the biomarkers alone (78, 83). Additionally, when CA125 was determined in combination with KLK6 and 13 instead of KLK10, it was found that the combination of these three biomarkers is a more sensitive test to detect early stage ovarian cancer than by determination of CA125 alone (40). Different from that approach, Rosen et al. determined KLK6 and 10 in sera of ovarian cancer patients lacking CA125 expression and reported that these

markers may have the potential to serve as sensitive markers for early detection of ovarian cancer (37).

Apart from their prognostic value, expression of several of the KLKs in tumour tissues or release into the blood/ascites is also measured to foresee response to adjuvant cancer therapy. For instance, for a cohort of ovarian cancer patients with recurrent disease, KLK4 protein expression as assessed by immunohistochemistry was determined to be elevated in tumour tissues of the group of patients progressing under treatment with the chemotherapeutic drug taxane (paclitaxel), but not in the group of responders (85). Likewise, there is a direct correlation between tumour tissue-associated elevation of KLK7 protein and failure of the ovarian cancer patients to respond to carboplatin ± taxol chemotherapy (86). Zheng et al. (55) combined determination of KLK6, 8, and 13 with cancer stage and debulking status to distinguish between responder (complete or partial) and non-responder ovarian cancer patients treated with platinum-based chemotherapy. It was also found that KLK10 expression is a predictive marker related to resistance of ovarian cancer tissues to chemotherapy (87).

Oikonomopoulou et al. (81) determined serum protein levels of KLK5-8, 10, and 11, together with the serum markers CA125, B7-H4 (a member of the B7 family of immune costimulatory proteins), regenerating protein IV, and spondin-2 at baseline, or after the first chemotherapy cycle, to predict a patient's response to carboplatin and/or taxol-based chemotherapy. The authors reported that a range of serum biomarkers encompassing CA125 and KLK5 and 7, would predict response to chemotherapy. All of these cancer biomarkers examined, except KLK7 and regenerating protein IV, were also powerful predictors of time-to-progression among the chemotherapy responders (81). Taken as a whole, KLKs have recently emerged as novel, promising predictive factors in ovarian cancer. However, owing to the lack of sufficiently effective alternative therapy models to treat advanced ovarian cancer patients, this knowledge is of limited clinical use.

Clinical utility of KLKs in cancers of the male reproductive tract

KLKs in prostate cancer

Prostate cancer is the second most common cancer in men worldwide (3) and the leading type of cancer in men in the USA (1, 2).

Definitive diagnosis of prostate cancer is done by biopsy. Malignant transformation of the epithelial cells lining the secretory glands of the prostate may manifest in early non-invasive lesions including high-grade prostatic intraepithelial neoplasia (HGPIN) (88). At diagnosis, tumour stage, Gleason score (89), and PSA (prostate-specific antigen, also known as kallikrein-related peptidase 3, KLK3) are the most accepted important predictors of prognosis of prostate cancer.

These parameters influence treatment strategies, which may include active surveillance; or surgery with or without a combination of radiation, steroid hormone or chemotherapeutic interventions for those cancers that are deemed aggressive. The observed biological heterogeneity in prostate cancer is reflected by the range of

survival among men diagnosed with the disease, with small localized cancers having the generally high 5-year average survival of >85% in industrialised countries (3), while more aggressive cancers having a much worse prognosis.

As such, current research is focusing on identifying genomic, transcriptomic, or proteomic biomarkers that distinguish between indolent and aggressive cancers for which molecular pathway may reveal therapeutic targets. To date, molecular profiling approaches have identified several potential markers, including the frequent losses of NKX3.1 (8p21) and PTEN (10q23), gains of AR and fusion of ETS family transcription factor genes with androgen-responsive promoters, amplifications of MYC, and mutations of TP53, PTEN, AR, RB1 or APC (90-94); in addition to miRNAs as potential prognostic biomarkers (95).

Prostate cancer is relatively slow growing, thus allowing a sufficient time for cancer to be discovered before it becomes incurable, for instance by use of the non-invasive PSA (prostate-specific antigen) screening test (96-102). Yet, today, PSA's (KLK3) effectiveness for screening and diagnosis of prostate cancer is not that obvious anymore and therefore questioned by several clinicians (100). Current controversies and major doubts regarding this matter of debate are explained and discussed in further detail by Fillon et al. (101).

In the normal prostate, all of the KLKs are expressed at the mRNA level and, except for KLK8, at the protein level as well (8, 16). In prostate cancer, elevated levels of KLK2, 4, and 12-15 mRNA and/or protein have been reported; three of those (KLK2, 14, and 15) are associated with poor prognosis; KLK4 is associated with favourable prognosis. Decreased mRNA or protein levels of KLK2, 3, 5-7, 10, 11, 13, and 15 have been reported of which KLK3 and 15 are associated with poor prognosis while KLK5 and 11 expression is associated with a favourable prognosis (► Table 4) (15, 102, 103).

The regulation of KLK expression in prostate cancer has been extensively studied, with evidence showing the regulatory role of various steroid hormones (104). Indeed, KLK2 and 3 possess defined steroid hormone binding sites (41, 104) while KLK1 and 4 possess putative steroid binding elements (105); the remaining KLKs fail to possess such defined elements (41, 104). Epigenetic studies suggest a role for differential DNA-methylation in KLK regulation (106-109) influencing aberrant expression. More recently, non-coding mRNAs, such as miRNAs, were suggested to be involved in the regulation of certain KLK genes (110).

Prostate cancer screening remains controversial, because of the inability of any biomarker to be detected early enough with high specificity a significant proportion of dangerous tumours. Yet, also in combination with PSA, several promising novel blood-based biomarkers for prostate cancer diagnosis, monitoring, or prediction of the course of the disease and response to cancer therapy are under investigation. e. g. KLK2, urokinase plasminogen activator (uPA) and its receptor uPAR, transforming growth factor-beta 1 (TGF-β 1) as well as interleukin-6 (IL-6) and its receptor IL-6R (111).

Table 4: KLKs present in normal and tumour tissues of patients afflicted with prostate cancer.

	Normal		Cancerous		References
	mRNA	Protein	mRNA	Protein	
KLK1	High	High	ND	ND	8, 142, 163–169
KLK2	High	High	↑	↑↓	8, 96, 102, 163–173
KLK3	High	High	↓	↓	7, 8, 96, 100, 102, 142, 163–170, 172, 174–176
KLK4	Moderate	Low	↑	↑	8, 12, 14, 16, 62, 163–169, 177–184
KLK5	Low	Low	↓	ND	8, 16, 163–169, 185, 186
KLK6	Low	Present	ND	↓	8, 12, 14, 16, 163–165, 168, 169, 180, 181
KLK7	Moderate	Present	↓	↓	8, 16, 163–169, 187–189
KLK8	Moderate	Absent	ND	ND	8, 16, 163–165, 168, 169, 190
KLK9	Low	High	ND	ND	8, 163–169
KLK10	High	Present	↓	↓	8, 12–14, 16, 109, 163–165, 168, 169, 178, 180, 181
KLK11	High	High	↓	↓	8, 16, 163–169, 187, 191–195
KLK12	Moderate	Present	ND	↑	8, 12, 14, 16, 179–181, 196–198
KLK13	Low	Low	↑	↑↓	8, 12, 14, 16, 163–169, 178, 180, 181
KLK14	High	Low	↑	↑	8, 16, 123, 163–169, 199, 200
KLK15	High	Low	↑	↓	8, 103, 163–169

↑ = Increased, ↓ = decreased compared to normal tissue. ND = Not determined.

KLKs in testicular cancer

Testicular cancer is the most common among Caucasian men between age 15 and 35; while relatively uncommon in Asia and Africa. More than 95% of testicular cancers are germ cell tumours. Testicular cancer is very well treatable (surgery, radiotherapy, chemotherapy) with a cure rate of ~95% if it has not spread outside the testicle, due to its extreme sensitivity to cisplatin-based chemotherapy, even when metastatic (112, 113). Indeed, if it has metastasised to organs or lymph nodes, the five-year relative survival rate is still high with ~72%.

Although two highly specific and sensitive serum markers: α -fetoprotein (AFP) and human chorionic gonadotropin (hCG) are not recommended for the diagnosis of testicular cancer in the absence of histologic confirmation, they are useful markers in the management of testicular germ cell tumours (114, 115). In the last years, novel biomarkers, e.g. glypican 3, SALL4, OCT3/4, SOX2, SOX17, OCT3/4, NANOG HMGA1, HMGA2, PATZ1, GPR30, and Aurora B were reported to discriminate between testicular cancer subgroups (116–118).

In the normal testis, all of the 15 KLKs are expressed at the mRNA level (8). Compared to other normal tissues, KLK proteins are only weakly/moderately expressed in testicular tissues; KLK15 is not expressed (8, 16) (► Table 5). Some of the KLKs have been shown to be of clinical value as biomarkers in testicular cancer as well, such as KLK5, 10, 11, 13, and 14 (119–124). mRNA expression data of these five imply, compared with normal testicular tissue, that these KLKs are downregulated in testicular cancer (119). Among those, lower mRNA levels in late stage (II/III) versus early stage (I) carcinomas indicate that KLK5 might be a marker

indicating a favourable prognosis (120). This might apply to the other four KLKs tested as well; even a possible tumour suppressive role of these KLKs in testicular cancer was discussed (121–123). In one of the studies, five splice variants of KLK13 were identified, exclusively expressed in normal testicular tissue, which are not present in the malignant counterpart (122). Full-length KLK13, on the other hand, is expressed both in the normal and the cancerous testicular tissue. To date, no study results relating to testicular cancer mRNA expression have been presented for the other ten KLKs; and at present, no results are available relating to the testicular tumour protein levels of KLK1–9 and 11–15.

KLKs in kidney cancer

Renal cell carcinoma (RCC), with an estimated incidence of 4% (64,770 of 1,638,910) and estimated 2.4% (13,570 of 577,190) cancer-related deaths for the year 2012 (1, 2) in the USA, comprises a histopathologically heterogeneous group with the majority of cancers being of the clear-cell subtype (80%); while the remaining subtypes include the papillary and chromophobe RCCs (124, 125). Along with tumour stage, most RCCs are graded according to the Fuhrman nuclear grading system (126). Surgical resection is the only known effective treatment for localised RCC. The most common molecular change in RCC is the loss of the VHL (von-Hippel-Lindau) gene function by deletion (127), epigenetic silencing (128), or protein regulation mediated by miRNAs (129). These findings have enabled the use of therapeutic agents that affect pathways including those of the VEGF pathway and TORC1 complex. Recent genomic and transcriptomic profiling (130) have

	Normal		Cancerous		References
	mRNA	Protein	mRNA	Protein	
KLK1	Moderate	Low	ND	ND	8, 41
KLK2	Moderate	Low	ND	ND	8
KLK3	Moderate	Moderate	ND	ND	8
KLK4	High	Low	ND	ND	8, 41
KLK5	High	Low	↓	ND	8, 16, 41, 119, 120, 163, 164
KLK6	High	Low	ND	ND	8, 16, 41, 164
KLK7	High	Low	ND	ND	8, 16, 41, 166
KLK8	High	Low	ND	ND	8, 190
KLK9	Low	Moderate	ND	ND	8, 41
KLK10	High	Low	↓	Expressed	8, 16, 41, 119, 121, 163
KLK11	High	Low	↓	ND	8, 16, 41, 163
KLK12	Low	Present	ND	ND	8, 16, 41
KLK13	Moderate	Low	↓	ND	8, 16, 41, 119, 122, 164
KLK14	High	Present	↓	ND	8, 16, 41, 119, 123
KLK15	Moderate	Absent	ND	ND	8, 41, 163

↑ = Increased, ↓ = decreased compared to normal tissue. ND = Not determined.

Table 5: KLKs present in normal and tumour tissues of patients afflicted with testicular/seminal cancer.

identified a number of recurrent copy-number changes including the gains/amplifications of genomic loci harbouring putative oncogenes such as MCM2, EGFR, and SEC61G; and losses of genomic regions harboring putative tumour suppressor genes including KIF1B, ARID1A, NHEJ1, KU80, and IGFBP5, among

others. Molecular profiling approaches have a great potential for improving RCC patient management (131). Moreover, recent transcriptomic profiling has identified two putative molecular subtypes of clear cell RCC with implications for outcome (132), and enabled accurate molecular classification of RCC subtypes (133).

	Normal		Cancerous		References
	mRNA	Protein	mRNA	Protein	
KLK1	High	Moderate	↓	↓	8, 16, 18, 41, 129, 163, 164, 201, 202
KLK2	Low	Low	Absent	ND	8, 129, 202
KLK3	Low	Mod	↓	ND	8, 129, 202
KLK4	Low	Low	Absent	ND	8, 129, 164
KLK5	Low	Mod	↓	↓	8, 16, 129, 134, 164
KLK6	High	Moderate/High	↓	↓	8, 12, 16, 18, 41, 129, 134, 152, 163, 164
KLK7	High	Low	↓	↓	8, 16, 18, 41, 129, 164, 166
KLK8	Moderate	Low	Absent	ND	8, 129, 190, 203
KLK9	Low	Mod	↓	ND	8, 129
KLK10	Low	High	Absent	↓	8, 13, 16, 129, 134, 164
KLK11	Low	Low	Absent	↓	8, 16, 129, 134, 164
KLK12	Low	Low	Absent	ND	8, 16, 129, 164
KLK13	Low	Low	↓	ND	8, 16, 129
KLK14	Moderate	Low	↓	ND	8, 16, 41, 129
KLK15	Low	Low	Absent	↓	8, 18, 41, 129

↑ = Increased, ↓ = decreased compared to normal tissue. ND = Not determined.

Table 6: KLKs present in normal and tumour tissues of patients afflicted with kidney cancer.

The normal kidney expresses all of the 15 KLKs, both at the mRNA and protein level (8, 16) (► Table 6). Petraki et al. (16) showed by immunohistochemistry that the eight KLKs they investigated (KLK5-7, and 10-14) are expressed by the epithelium of the urinary tubuli of the kidney while glomeruli are negative. In RCC, there is a consistent decrease (KLK1, 3, 5-7, 9-11, and 13-15) or absence (KLK2, 4, 8, 10-12, and 15) of the KLKs at either the mRNA and/or protein levels (16, 18, 129, 134). Additionally, studies by Gabril et al. (18) revealed differential immunostaining pattern of KLKs in RCC when compared to normal kidney tissue. KLKs were also found to be differentially expressed in the different RCC subtypes. There was an overall decrease in KLK1 expression in clear cell RCC compared to adjacent normal kidney; although relatively stronger KLK1 immunoreactivity was seen in high-grade compared with low-grade tumours. These findings were supported by those of White et al. (129), showing that the decreased expression of KLK1 and 3 was significantly associated with the clear cell subtype.

KLK1 showed weak expression in papillary RCC and negative to weak focal expression in chromophobe RCC and oncocytomas. In clear cell RCC, there was decreased KLK6 expression in higher grade compared with lower grade cancers. Papillary RCC showed strong diffuse KLK6 cytoplasmic protein expression with focal apical accentuation. In contrast, oncocytoma and chromophobe RCC showed no expression in tumour cells with only KLK6 expression in the blood vessels of the stroma. For KLK7, there was decreased protein expression observed in high-grade compared to low-grade-tumours. Oncocytomas showed diffuse, strong granular cytoplasmic expression of KLK7 protein while chromophobe RCC showed focal weak homogeneous cytoplasmic staining. Papillary RCC showed strong, apical cytoplasmic KLK7 protein staining. For KLK15 protein, clear cell RCCs showed cytoplasmic staining, weak to negative expression in papillary RCCs, and no expression in chromophobe RCC, urothelial carcinomas, or oncocytomas. Though still in its early stages, these findings indicate a good potential for the use of KLKs as adjuvant tissue markers to distinguish RCC subtypes.

Most of the potential clinical utility of KLKs in RCC is related to their prognostic value and their ability to distinguish between the RCC subtypes. KLK5, 6, 10, and 11 were assessed for their potential to predict the course of the RCC disease (134). It was only that KLK6 was of prognostic value to predict disease-specific survival; KLK5, 10, and 11 were of no effect (134).

KLK expression may be regulated transcriptionally and/or post-transcriptionally (102, 105, 106, 135). White et al. (129) demonstrated that the differential mRNA expression of KLKs in RCC may be mediated by the aberrant expression of miRNAs. Specifically, luciferase assays demonstrated the ability of hsa-let-7f to target the 3'untranslated region (UTR) of KLK10 in the RCC cell line ACHN, while transient transfection of miR-224 to HEK-293 cells resulted in the decrease of *KLK1* mRNA expression.

Table 7: KLKs present in normal and tumour tissues of patients afflicted with bladder cancer.

	Normal ^a		Cancerous		References
	mRNA	Protein	mRNA	Protein	
KLK1	ND	Low	ND	ND	8
KLK2	ND	Absent	ND	ND	8
KLK3	ND	Absent	ND	Absent	8, 139, 140–142
KLK4	ND	Absent	ND	ND	8
KLK5	ND	Absent	↑	↑	8, 16, 138
KLK6	ND	Absent	↑	↑	8, 13, 16, 138
KLK7	ND	Low	ND	ND	8
KLK8	ND	Absent	↑	ND	8, 138
KLK9	ND	Moderate	↑	ND	8, 138
KLK10	ND	Absent	ND	↑	8, 12, 13, 16
KLK11	ND	Absent	ND	↑	8, 16
KLK12	ND	Absent	ND	ND	8
KLK13	ND	Low	ND	ND	8
KLK14	ND	Moderate	ND	ND	8
KLK15	ND	Absent	ND	ND	8

↑ = Increased, ↓ = decreased compared to normal tissue. ND = Not determined.

KLKs in urinary bladder cancer

Urinary bladder carcinomas rank ninth in overall cancer incidence worldwide and are found more commonly among men than women. The vast majority of bladder cancers are of the urothelial type, superficial, and are typically treated by transurethral resection or intravesical immunotherapy and chemotherapy (136, 137). For these patients, the 5-year survival rate is nearly 90%. However, approximately 50-70% of tumours recur with some evolving into a more invasive phenotype. Patients who present with the infiltrative type have only a 60% 5-year survival (136, 137).

Approximately 10-15% of the non-invasive lesions progress into the muscle invasive-disease. Molecular genetics and cytogenetics have identified alterations of chromosome 9 to the normal urothelium as the earliest change prior to the divergence into the two different pathways. Disruptions of the PI3K-AKT axis and alterations of HRAS are generally associated with the non-invasive subtype; whereas alterations in genes affecting cell cycle and apoptosis include *TP53*, *CDKN2A*, *CCND1*, *CDKN1B*, or *RB*.

To date there is relatively little known about the expression of KLKs in the normal urinary bladder or in bladder cancer and their potential use for clinical management of that malignancy. No information is available in the scientific literature concerning the mRNA expression levels in the normal bladder of any of the KLKs. Regarding the KLK protein expression in the urinary bladder, KLK1, 7, 9, 13, and 14 are expressed, the other 10 KLKs are absent (► Table 7). For most of the KLKs, their expression levels in uri-

nary bladder carcinoma were not investigated, except for *KLK5*, 6, 8, and 9 *KLK mRNA* and *KLK5*, 6, 10, and 11 protein, which are all increased compared to the normal tissue counterpart (16, 18, 138). Other studies have focused on the putative clinical utility of *KLK3* in bladder cancer. These studies have shown and confirmed the lack of protein expression of *KLK3* in most of the bladder cancers investigated (139-141).

Perspective

This review summarises the clinical importance of various KLKs in several types of female and male urogenital malignancies, to give an overview and interpretation of the clinical impact of certain KLKs in these types of cancers. Although KLKs are primarily known for their biomarker value in prostate, ovarian, breast, and gastrointestinal cancers, regarding prediction of the course of the disease and response to cancer therapy, more recent data point to an important role of certain KLKs in other malignancies as well, such as those of the lung, brain, head and neck, the kidney, the endometrium, cervix, testis, and the urinary bladder (7, 15, 204).




Figure 1: Comparative mRNA (A) and protein (B) expression in normal tissues of the urogenital tract.




Figure 2: Comparative mRNA (A) and protein (B) expression in cancerous tissues of the urogenital tract.

KLKs, which represent a major tissue-associated proteolytic system, stand for a rich source of biomarkers that allow molecular classification, early diagnosis and prognosis of human malignancies as well as prediction of response or failure to cancer-directed drugs (7, 15, 204, 205). Without a doubt, these data point to the important role of various KLKs for improving individualized cancer therapy and thus clinical management of the cancer patients, especially of those patients who are likely to respond, while directing those who would not benefit towards alternative therapeutic options.

Regarding KLK expression in female and male urogenital organs and their tumour-affected counterparts, a large number of the KLKs are expressed in the normal, unaffected organs, at the mRNA and/or protein level (►Figure 1), with KLK1 and 14 expressed in all of the organs listed. Remarkably, all of the KLKs are expressed in the normal prostate, testis, and kidney whereas the uterus, the ovary, and the urinary bladder are expressing a limited number of KLKs only.

Regarding the KLK expression levels present in malignant urogenital tumours, remarkably, expression data for KLK1 (also known as tissue kallikrein) have been published only for endometrium and kidney cancer. Most of the information regarding KLK expression in tumour-affected organs is available for ovarian cancer; all of the 12 KLKs tested were found to be elevated in the malignant state, depicting them as valuable biomarkers to distinguish between the normal and the cancerous phenotype. Different from that, in kidney cancer, a series of KLKs were found to be downregulated, other KLKs were not expressed (►Figure 2).

Evidently, depending on the type of cancer or cancer stage, individual KLKs may show characteristics of a Janus-faced behaviour, by either expanding or inhibiting cancer progression and metastasis. Therefore, in order to interpret the clinical impact of different KLKs on the broad range of malignancies, the exact mechanisms of action of KLKs in distinct tumour entities and their microenvironments will need to be further elucidated.

KLKs often work in cascades by activating each other or function in signaling pathways involving kinases, phosphatases, and other proteases such as the urokinase-type plasminogen activator (uPA) (206). At first glance, the KLK proteases are characterised by high sequence similarities, which may pose many challenges impacting the development of highly specific inhibitors. Yet, KLKs do show significant differences in their substrate specificities, which have facilitated development of targeted KLK inhibitors (207). Some of the novel KLK-directed therapeutics have been tested in experimental animal models; two, affecting KLK3 (PSA) are already in phase II/III clinical trials (205).

Conflict of interest

None declared.

References

1. Siegel R, DeSantis C, Virgo K, et al. Cancer treatment and survivorship statistics, 2012. *CA Cancer J Clin* 2012; 62: 220-241.

2. Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin* 2012; 62: 10-29.
3. GLOBOCAN 2008. Estimated cancer incidence, mortality, prevalence and disability-adjusted life years (DALYs) worldwide in 2008. Available at: <http://globocan.iarc.fr/>.
4. FIGO, International Federation of gynaecology and Obstetrics. Available at: <http://www.figo.org>.
5. National Cancer Institute. Available at: <http://www.cancer.gov/dictionary/?Cdrid=45618>.
6. Schmitt M, Magdolen V. Using kallikrein-related peptidases (KLK) as novel cancer biomarkers. *Thromb Haemost* 2009; 101: 222-224.
7. Avgeris M, Mavridis K, Scorilas A. Kallikrein-related peptidase genes as promising biomarkers for prognosis and monitoring of human malignancies. *Biol Chem* 2010; 391: 505-511.
8. Shaw JL, Diamandis EP. Distribution of 15 human kallikreins in tissues and biological fluids. *Clin Chem* 2007; 53: 1423-1432.
9. Shaw JL, Petraki C, Watson C, et al. Role of tissue kallikrein-related peptidases in cervical mucus remodeling and host defense. *Biol Chem* 2008; 389: 1513-1522.
10. Jin H, Nagai N, Shigemasa K, et al. Expression of tumour-associated differentially expressed Gene-14 (TADG-14/KLK8) and its protein hK8 in uterine endometria and endometrial carcinomas. *Tumour Biol* 2006; 27: 274-282.
11. Borgho CA, Grass L, Soosaipillai A, et al. Human kallikrein 14: a new potential biomarker for ovarian and breast cancer. *Cancer Res* 2003; 24: 9032-9041.
12. Petraki CD, Karavana VN, Skoufogiannis PT, et al. The spectrum of human kallikrein 6 (zyme/protease M/neurosin) expression in human tissues, as assessed by immunohistochemistry. *J Histochem Cytochem* 2001; 49: 1431-1441.
13. Petraki CD, Karavana VN, Luo LY, et al. Human kallikrein 10 expression in normal tissues by immunohistochemistry. *J Histochem Cytochem* 2002; 50: 1247-1261.
14. Petraki CD, Karavana VN, Diamandis EP. Human kallikrein 13 expression in normal tissues. An immunohistochemical study. *J Histochem. Cytochem* 2003; 51: 493-501.
15. Avgeris M, Mavridis K, Scorilas A. Kallikrein-related peptidases in prostate, breast, and ovarian cancers: from pathobiology to clinical relevance. *Biol Chem* 2012; 393: 301-317.
16. Petraki CD, Papanastasiou PA, Karavana VN, et al. Cellular distribution of human tissue kallikreins: immunohistochemical localisation. *Biol Chem* 2006; 387: 653-663.
17. Xi Z, Klok T.I, Korkmaz K, et al. Kallikrein 4 is a predominantly nuclear protein and is overexpressed in prostate cancer. *Cancer Res* 2004; 64: 2365-2370.
18. Gabril M, White NM, Moussa M, et al. Immunohistochemical analysis of kallikrein-related peptidases in the normal kidney and renal tumours: potential clinical implications. *Biol Chem* 2010; 391: 403-409.
19. Emami N, Diamandis EP. Utility of kallikrein-related peptidases (KLKs) as cancer biomarkers. *Clin Chem* 2008; 54: 1600-1607.
20. Bittoni MA, Fisher JL, Fowler JM, et al. Assessment of the effects of severe obesity and lifestyle risk factors on stage of endometrial cancer. *Cancer Epidemiol Biomarkers Prev* 2013; 1: 76-81.
21. Folkens AK, Longare TA. Hereditary gynaecological malignancies: advances in screening and treatment. *Histopathology* 2013; 1: 2-30.
22. Bovicelli A, D'Andrilli G, Giordano A, et al. Conservative treatment of early endometrial cancer. *J Cell Physiology* 2012 Epub ahead of print.
23. Sood AK, Buller RE, Burger RA, et al. Value of preoperative CA 125 level in the management of uterine cancer and prediction of clinical outcome. *Obstet gynaecol* 1997; 3: 441-447.
24. Myers SA, Clements JA. Kallikrein 4 (KLK4), a new member of the human kallikrein gene family is up-regulated by estrogen and progesterone in the human endometrial cancer cell line, KLE. *J Clin Endocrinol Metab* 2001; 5: 2323-2326.
25. Zhang SQ, Cai B, Liu L, et al. Kallikrein 4 overexpression in endometrial carcinoma and upregulation by estrogen via mitogen-activated protein kinase signal pathway. *Int J gynaecol Cancer* 2009; 8: 1377-1383.
26. Santin AD, Diamandis EP, Bellone S, et al. Overexpression of kallikrein 10 (hK10) in uterine serous papillary carcinomas. *Am J Obstet gynaecol* 2006; 5: 1296-1302.
27. Santin AD, Diamandis EP, Bellone S, et al. Human kallikrein 6: a new potential serum biomarker for uterine serous papillary cancer. *Clin Cancer Res* 2005; 9: 3320-3325.

28. zur Hausen H. Papillomaviruses—to vaccination and beyond. *Biochemistry* 2008; 5: 498-503.
29. Rose PG, Bundy BN, Watkins EB, et al. Concurrent cisplatin-based radiotherapy and chemotherapy for locally advanced cervical cancer. *N Engl J Med* 1999; 15: 1144-1153.
30. Micke O, Bruns F, Schäfer U, et al. The impact of squamous cell carcinoma (SCC) antigen in patients with advanced cancer of uterine cervix treated with (chemo)-radiotherapy. *Anticancer Res* 2005; 3A: 1663-1666.
31. Santin AD, Cane S, Bellone S, et al. The serine protease stratum corneum chymotryptic enzyme (kallikrein 7) is highly overexpressed in squamous cervical cancer cells. *gynaecol Oncol* 2004; 2: 283-288.
32. Termini L, Maciag PC, Soares FA, et al. Analysis of human kallikrein 7 expression as a potential biomarker in cervical neoplasia. *Int J Cancer* 2010; 2: 485-490.
33. Tian X, Shigemasa K, Hirata E, et al. Expression of human kallikrein 7 (hK7/SCCE) and its inhibitor antileukoprotease (ALP/SLPI) in uterine endocervical glands and in cervical adenocarcinomas. *Oncol Rep* 2004; 5: 1001-1006.
34. Cane S, Bignotti E, Bellone S, et al. The novel serine protease tumour-associated differentially expressed gene-14 (KLK8/Neurospine/Ovasin) is highly overexpressed in cervical cancer. *Am J Obstet gynaecol* 2004; 1: 60-66.
35. Lengyel E. Ovarian cancer development and metastasis. *Am J Pathol* 2010; 3: 1053-1064.
36. Yip P, Chen TH, Seshiah P, et al. Comprehensive serum profiling for the discovery of epithelial ovarian cancer biomarkers. *PLoS One* 2011; 12: e29533.
37. Rosen DG, Wang L, Atkinson JN, et al. Potential markers that complement expression of CA125 in epithelial ovarian cancer. *gynaecol Oncol* 2005; 2: 267-277.
38. Dorn J, Schmitt M, Kates R, et al. Primary tumour levels of human tissue kallikreins affect surgical success and survival in ovarian cancer patients. *Clin Cancer Res* 2007; 6: 1742-1748.
39. Yousef GM, Diamandis EP. The human kallikrein gene family: new biomarkers for ovarian cancer. *Cancer Treat Res* 2009; 149: 165-187.
40. White N, Mathews M, Yousef GM, et al. Human kallikrein related peptidases 6 and 13 in combination with CA125 is a more sensitive test for ovarian cancer than CA125 alone. *Cancer Biomark* 2009; 6: 279-287.
41. Yousef GM, Diamandis EP. The new human tissue kallikrein gene family: structure function and association to disease. *Endocr Rev* 2001; 22: 184-204.
42. Bignotti E, Tassi RA, Calza S, et al. Differential gene expression profiles between tumour biopsies and short-term primary cultures of ovarian serous carcinomas: identification of novel molecular biomarkers for early diagnosis and therapy. *gynaecol Oncol* 2006; 2: 405-416.
43. Dong Y, Kaushal A, Bui L, et al. Human kallikrein 4 (KLK4) is highly expressed in serous ovarian carcinomas. *Clin Cancer Res* 2001; 8: 2363-2371.
44. Yousef GM, Polymeris ME, Yacoub GM, et al. Parallel overexpression of seven kallikrein genes in ovarian cancer. *Cancer Res* 2003; 9: 2223-2227.
45. Adib TR, Henderson S, Perrett C, et al. Predicting biomarkers for ovarian cancer using gene-expression microarrays. *Br J Cancer* 2004; 3: 686-692.
46. Tanimoto H, Underwood LJ, Shigemasa K, et al. Increased expression of protease M in ovarian tumours. *Tumour Biol* 2001; 1: 11-18.
47. Tanimoto H, Underwood LJ, Shigemasa K, et al. The stratum corneum chymotryptic enzyme that mediates shedding and desquamation of skin cells is highly overexpressed in ovarian tumour cells. *Cancer* 1999; 10: 2047-2082.
48. Dong Y, Kaushal A, Brattsand M, et al. Differential splicing of KLK5 and KLK7 in epithelial ovarian cancer produces novel variants with potential as cancer biomarkers. *Clin Cancer Res* 2003; 5: 1710-1720.
49. Shvartsman HS, Lu KH, Lee J, et al. Overexpression of kallikrein 10 in epithelial ovarian carcinomas. *gynaecol Oncol* 2003; 1: 44-50.
50. Gilks CB, Vancerhyden BC, Zhu S, et al. Distinction between serous tumours of low malignant potential and serous carcinomas based on global mRNA expression profiling. *gynaecol Oncol* 2005; 3: 684-694.
51. Yousef GM, Fraccioli S, Scorilas A, et al. Steroid hormone regulation and prognostic value of the human kallikrein gene 14 in ovarian cancer. *Am J Clin Pathol* 2003; 3: 346-355.
52. Yousef GM, Scorilas A, Katsaros D, et al. Prognostic value of the human kallikrein gene 15 expression in ovarian cancer. *J Clin Oncol* 2003; 16: 3119-3126.
53. Shan SJ, Scorilas A, Katsaros D, et al. Unfavourable prognostic value of human kallikrein 7 quantified by ELISA in ovarian cancer cytosols. *Clin Chem* 2006; 10: 1879-1886.
54. Dorn J, Harbeck N, Kates R, et al. Disease processes may be reflected by correlations among tissue kallikrein proteases but not with proteolytic factors uPA and PAI-1 in primary ovarian carcinoma. *Biol Chem* 2006; 8: 1121-1128.
55. Zheng Y, Katsaros D, Shan SJ, et al. A multiparametric panel for ovarian cancer diagnosis, prognosis, and response to chemotherapy. *Clin Cancer Res* 2007; 23: 6984-6992.
56. Dorn J, Milou V, Kulasingam V, et al. Clinical relevance of kallikrein-related peptidases in ovarian cancer. In: *Kallikrein-related peptidases. Volume 2 – Novel cancer-related biomarkers*. deGruyter, Berlin, 2012. pp. 145-166.
57. O'Mara TA, Nagle CM, Batra J, et al. Kallikrein-related peptidase 3 KLK3/PSA single nucleotide polymorphisms and ovarian cancer survival. *Twin Res Hum Genet* 2011; 14: 323-327.
58. Obiezu CV, Scorilas A, Katsaros K, et al. Higher human kallikrein gene 4 (KLK4) expression indicates poor prognosis of ovarian cancer patients. *Clin Cancer Res* 2001; 8: 2380-2386.
59. Diamandis EP, Borgoño CA, Scorilas A, et al. Immunofluorometric quantification of human kallikrein 5 expression in ovarian cancer cytosols and its association with unfavourable patient prognosis. *Tumour Biol* 2003; 6: 299-309.
60. Kim H, Scorilas A, Katsaros D, et al. Human kallikrein gene 5 (KLK5) expression is an indicator of poor prognosis in ovarian cancer. *Br J Cancer* 2001; 5: 643-650.
61. Hoffman BR, Katsaros D, Scorilas A, et al. Immunofluorometric quantitation and histochemical localisation of kallikrein 6 protein in ovarian cancer tissue: a new independent unfavourable prognostic biomarker. *Br J Cancer* 2002; 7: 763-771.
62. Seiz L, Dorn J, Kotsch M, et al. Stromal cell-associated expression of kallikrein-related peptidase 6 (KLK6) indicates poor prognosis of ovarian cancer patients. *Biol Chem* 2012; 391: 391-401.
63. Kyriakopoulou LG, Yousef GM, Scorilas A, et al. Prognostic value of quantitatively assessed KLK7 expression in ovarian cancer. *Clin Biochem* 2003; 36: 135-143.
64. Yousef GM, Kyriakopoulou LG, Scorilas A, et al. Quantitative expression of the human kallikrein gene 9 (KLK9) in ovarian cancer: a new independent and favourable prognostic marker. *Cancer Res* 2001; 21: 7811-7818.
65. Borgoño CA, Kishi T, Scorilas A, et al. Human kallikrein 8 protein is a favourable prognostic marker in ovarian cancer. *Clin Cancer Res* 2006; 5: 1487-1493.
66. Kishi T, Grass L, Soosaipillai A, et al. Human kallikrein 8 a novel biomarker for ovarian carcinoma. *Cancer Res* 2003; 63: 2771-2774.
67. Shigemasa K, Tian X, Gu L, et al. Human kallikrein 8 (hK8/TADG-14) expression is associated with an early clinical stage and favourable prognosis in ovarian cancer. *Oncol Rep* 2004; 11: 1153-1159.
68. Magklara A, Scorilas A, Katsaros D, et al. The human KLK8 neurospine/ovasin gene: identification of two novel splice variants and its prognostic value in ovarian cancer. *Clin Cancer Res* 2001; 7: 806-811.
69. Kountourakis P, Psyrri A, Scorilas A, et al. Expression and prognostic significance of kallikrein-related peptidase 8 protein levels in advanced ovarian cancer by using automated quantitative analysis. *Thromb Haemost* 2009; 101: 541-546.
70. Luo LY, Katsaros D, Scorilas A, et al. Prognostic value of human kallikrein 10 expression in epithelial ovarian carcinoma. *Clin Cancer Res* 2001; 8: 2372-2379.
71. Borgoño CA, Fraccioli S, Yousef GM, et al. Favourable prognostic value of tissue human kallikrein 11 (hK11) in patients with ovarian carcinoma. *Int J Cancer* 2003; 4: 605-610.
72. Diamandis EP, Borgoño CA, Scorilas A, et al. Human kallikrein 11: an indicator of favourable prognosis in ovarian cancer patients. *Clin Biochem* 2004; 9: 823-829.
73. Scorilas A, Borgoño CA, Harbeck N, et al. Human kallikrein 13 protein in ovarian cancer cytosols: a new favourable prognostic marker. *J Clin Oncol* 2004; 4: 768-785.
74. Shigemasa K, Gu L, Tanimoto H, et al. Human kallikrein gene 11 (KLK11) mRNA overexpression is associated with poor prognosis in patients with epithelial ovarian cancer. *Clin Cancer Res* 2004; 8: 2766-2770.
75. White NM, Mathews M, Yousef GM, et al. KLK6 and KLK13 predict tumour recurrence in epithelial ovarian carcinoma. *Br J Cancer* 2009; 101: 1107-1113.
76. Dorn J, Magdolen V, Gkazepis A, et al. Circulating biomarker tissue kallikrein-related peptidase KLK5 impacts ovarian cancer patients' survival. *Ann Oncol* 2011; 8: 1783-1790.
77. Bandiera E, Zanotti L, Bignotti E, et al. Human kallikrein 5: an interesting novel biomarker in ovarian cancer patients that elicits humoral response. *Int J gynaecol Cancer* 2009; 6: 1015-1021.

78. Diamandis EP, Scorilas A, Fraccioli S, et al. Human kallikrein 6 (hK6): a new potential serum biomarker for diagnosis and prognosis of ovarian carcinoma. *J Clin Oncol* 2003; 6: 1035-1043.
79. Koh SC, Razvi K, Chan YH, et al. The association with age, human tissue kallikreins 6 and 10 and hemostatic markers for survival outcome from epithelial ovarian cancer. *Arch gynaecol Obstet* 2011; 1: 183-190.
80. Luo LY, Katsaros D, Scorilas A, et al. The serum concentration of human kallikrein 10 represents a novel biomarker for ovarian cancer diagnosis and prognosis. *Cancer Res* 2003; 4: 807-811.
81. Oikonomopoulou K, Li L, Zheng Y, et al. Prediction of ovarian cancer prognosis and response to chemotherapy by a serum-based multiparametric biomarker panel. *Br J Cancer* 2008; 7: 1103-1113.
82. El Sherbini MA, Sallam MM, Shaban EA, et al. Diagnostic value of serum kallikrein-related peptidases 6 and 10 versus CA125 in ovarian cancer. *Int J gynaecol Cancer* 2011; 4: 625-632.
83. Koh SC, Huak CY, Lutan D, et al. Combined panel of serum human tissue kallikreins and CA-125 for the detection of epithelial ovarian cancer. *J gynaecol Oncol* 2012; 23: 175-181.
84. Shih IeM, Salani R, Fiegl M, et al. *gynaecol Oncol* 2007; 105: 501-507.
85. Xi Z, Kaern J, Davidson B, et al. Kallikrein 4 is associated with paclitaxel resistance in ovarian cancer. *gynaecol Oncol* 2004; 1: 80-85.
86. Dong Y, Tan OL, Loessner D, et al. Kallikrein-related peptidase 7 promotes multicellular aggregation via the alpha(5)beta(1) integrin pathway and paclitaxel chemoresistance in serous epithelial ovarian carcinoma. *Cancer Res* 2010; 70: 2624-2633.
87. Cheng L, Lu W, Kulkarni B, et al. Analysis of chemotherapy response programs in ovarian cancers by the next-generation sequencing technologies. *gynaecol Oncol* 2010; 117: 159-169.
88. Xin L. 2012; Cells of origin for cancer: an updated view from prostate cancer. *Oncogene* 2012; Epub ahead of print.
89. Epstein JI, Allsbrook WC, Jr, Amin MB, et al. Update on the Gleason grading system for prostate cancer: results of an international consensus conference of urologic pathologists. *Adv Anat Pathol* 2006; 13: 57-59.
90. Barbieri CE, Baca SC, Lawrence MS, et al. Exome sequencing identifies recurrent SPOP FOXA1 and MED12 mutations in prostate cancer. *Nat Genet* 2012; 44: 685-689.
91. Berger MF, Lawrence MS, Demicheli F, et al. The genomic complexity of primary human prostate cancer. *Nature* 2011; 470: 214-220.
92. Beroukhi R, Mermel CH, Porter D, et al. The landscape of somatic copy-number alteration across human cancers. *Nature* 2010; 463: 899-905.
93. Grasso CS, Wu YM, Robinson DR, et al. The mutational landscape of lethal castration-resistant prostate cancer. *Nature* 2012; 487: 239-243.
94. Taylor BS, Schultz N, Hieronymus H, et al. Integrative genomic profiling of human prostate cancer. *Cancer Cell* 2010; 18: 11-22.
95. Fendler A, Jung M, Stephan C, et al. miRNAs can predict prostate cancer biochemical relapse and are involved in tumour progression. *Int J Oncol* 2011; 39: 1183-1192.
96. Becker C, Noldus J, Diamandis E, et al. The role of molecular forms of prostate-specific antigen (PSA or hK3) and of human glandular kallikrein 2 (hK2) in the diagnosis and monitoring of prostate cancer and in extra-prostatic disease. *Crit Rev Clin Lab Sci* 2001; 38: 357-399.
97. Kuriyama M, Wang MC, Papsidero LD, et al. Quantitation of prostate-specific antigen in serum by a sensitive enzyme immunoassay. *Cancer Res* 1980; 40: 4658-4662.
98. Lilja H, Ulmert D, Vickers AJ. Prostate-specific antigen and prostate cancer: prediction detection and monitoring. *Nat Rev Cancer* 2008; 8: 268-278.
99. Stamey TA, Yang N, Hay AR, et al. Prostate-specific antigen as a serum marker for adenocarcinoma of the prostate. *N Engl J Med* 1987; 317: 909-916.
100. Ulmert D, O'Brien MF, Bjartell AS, et al. Prostate kallikrein markers in diagnosis risk stratification and prognosis. *Nat Rev Urol* 2009; 6: 384-391.
101. Fillon M. Urology meeting highlights prostate, bladder cancers. *J Natl Cancer Inst* 2012; 104: 1348-1349.
102. Nam RK, Diamandis EP, Toi A, et al. Serum human glandular kallikrein-2 protease levels predict the presence of prostate cancer among men with elevated prostate-specific antigen. *J Clin Oncol* 2000; 18: 1036-1042.
103. Mavridis K, Avgeris M, Koutalellis G, et al. Expression analysis and study of the KLK15 mRNA splice variants in prostate cancer and benign prostatic hyperplasia. *Cancer Sci* 2010; 101: 693-699.
104. Lawrence MG, Lai J, Clements JA. Kallikreins on steroids: structure and hormonal regulation of prostate-specific antigen and the extended kallikrein locus. *Endocr Rev* 2010; 31: 407-446.
105. Lai J, Myers SA, Lawrence MG, et al. Direct progesterone receptor and indirect androgen receptor interactions with the kallikrein-related peptidase 4 gene promoter in breast and prostate cancer. *Mol Cancer Res* 2009; 7: 129-141.
106. Olkhov-Mitsel E, van der Kwast T, Kron KJ, et al. Quantitative DNA methylation analysis of genes coding for kallikrein-related peptidases 6 and 10 as biomarkers for prostate cancer. *Epigenetics* 2012; 7: 1037-1045.
107. Pampalakis G, Diamandis EP, Sotiropoulou G. The epigenetic basis for the aberrant expression of kallikreins in human cancers. *Biol Chem* 2006; 387: 795-799.
108. Pasic MD, Olkhov E, Bapat B, et al. Epigenetic regulation of kallikrein-related peptidases: there is a whole new world out there. *Biol Chem* 2012; 393: 319-330.
109. Sidiropoulos M, Pampalakis G, Sotiropoulou G, et al. Downregulation of human kallikrein 10 (KLK10/NES1) by CpG island hypermethylation in breast ovarian and prostate cancers. *Tumour Biol* 2005; 26: 324-336.
110. White NM, Youssef YM, Fendler A, et al. The miRNA-kallikrein axis of interaction: a new dimension in the pathogenesis of prostate cancer. *Biol Chem* 2012; 393: 379-389.
111. Shariat SF, Semjonow A, Lilja H, et al. Tumour markers in prostate cancer I: blood-based markers. *Acta Oncol* 2011; 50 (Suppl 1): 61-75.
112. Viatori M. Testicular cancer. *Semin Oncol Nurs* 2012; 28: 180-189.
113. Okamoto K. Epigenetics: a way to understand the origin and biology of testicular germ cell tumours. *Int J Urol* 2012; 19: 504-511.
114. Barlow LJ, Badalato GM, McKiernan JM. Serum tumour markers in the evaluation of male germ cell tumours. *Nat Rev Urol* 2010; 7: 610-617.
115. Salem M, Gilligan T. Serum tumour markers and their utilisation in the management of germ-cell tumours in adult males. *Expert Rev Anticancer Ther* 2011; 11: 1-4.
116. Emerson RE, Ulbright TM. Intratubular germ cell neoplasia of the testis and its associated cancers: the use of novel biomarkers. *Pathology* 2010; 42: 344-355.
117. Favilla V, Cimino S, Madonia M, Morgia G. New advances in clinical biomarkers in testis cancer. *Front Biosci* 2010; 2: 456-477.
118. Chieffo P. New prognostic markers and potential therapeutic targets in human testicular germ cell tumours. *Curr Med Chem* 2011; 18: 5033-5040.
119. Luo LY, Yousef G, Diamandis EP. Human tissue kallikreins and testicular cancer. *APMIS* 2003; 111: 225-232.
120. Yousef GM, Obiezu CV, Jung K, et al. Differential expression of Kallikrein gene 5 in cancerous and normal testicular tissues. *Urology* 2002; 60: 714-718.
121. Luo LY, Rajpert-de Meyts ER, Jung K, et al. Expression of the normal epithelial cell-specific 1 (NES1; KLK10) candidate tumour suppressor gene in normal and malignant testicular tissue. *Br J Cancer* 2001; 85: 220-224.
122. Chang A, Yousef GM, Jung K, et al. Identification and molecular characterization of five novel kallikrein gene 13 (KLK13; KLK-L4) splice variants: differential expression in the human testis and testicular cancer. *Anticancer Res* 2001; 21: 3147-3152.
123. Yousef GM, Magklara A, Chang A, et al. Cloning of a new member of the human kallikrein gene family, KLK14, which is down-regulated in different malignancies. *Cancer Res* 2001; 61: 3425-3431.
124. Arsanious A, Bjarnason GA, Yousef GM. From bench to bedside: current and future applications of molecular profiling in renal cell carcinoma. *Mol Cancer* 2009; 8: 20.
125. McLaughlin JK, Lipworth L, Tarone RE. Epidemiologic aspects of renal cell carcinoma. *Semin Oncol* 2006; 33: 527-533.
126. Fuhrman SA, Lasky LC, Limas C. Prognostic significance of morphologic parameters in renal cell carcinoma. *Am J Surg Pathol* 1982; 6: 655-663.
127. Latif F, Tory K, Gnarr J, et al. Identification of the von Hippel-Lindau disease tumour suppressor gene. *Science* 1993; 260: 1317-1320.
128. Moore LE, Nickerson ML, Brennan P, et al. von Hippel-Lindau (VHL) inactivation in sporadic clear cell renal cancer: associations with germline VHL polymorphisms and etiologic risk factors. *PLoS Genet* 2011; 7: e1002312.
129. White NM, Bui A, Mejia-Guerrero S, et al. Dysregulation of kallikrein-related peptidases in renal cell carcinoma: potential targets of miRNAs. *Biol Chem* 2010; 391: 411-423.
130. Girgis AH, Iakovlev VV, Beheshti B, et al. Multilevel whole-genome analysis reveals candidate biomarkers in clear cell renal cell carcinoma. *Cancer Res* 2012; 72: 5273-5284.
131. White NM, Yousef GM. Translating molecular signatures of renal cell carcinoma into clinical practice. *J Urol* 2011; 186: 9-11.

132. Beleut M, Zimmermann P, Baudis M, et al. Integrative genome-wide expression profiling identifies three distinct molecular subgroups of renal cell carcinoma with different patient outcome. *BMC Cancer* 2012; 12: 310.
133. Youssef YM, White NM, Grigull J, et al. Accurate molecular classification of kidney cancer subtypes using microRNA signature. *Eur Urol* 2011; 59: 721-730.
134. Petraki CD, Gregorakis AK, Vaslamatzis MM, et al. Prognostic implications of the immunohistochemical expression of human kallikreins 5, 6, 10 and 11 in renal cell carcinoma. *Tumour Biol* 2006; 27: 1-7.
135. Borgono CA, Diamandis EP. The emerging roles of human tissue kallikreins in cancer. *Nat Rev Cancer* 2004; 4: 876-890.
136. Donat SM. Evaluation and follow-up strategies for superficial bladder cancer. *Urol Clin North Am* 2003; 30: 765-776.
137. von der Maase H, Sengelov L, Roberts JT, et al. Long-term survival results of a randomized trial comparing gemcitabine plus cisplatin with methotrexate vinblastine doxorubicin plus cisplatin in patients with bladder cancer. *J Clin Oncol* 2005; 23: 4602-4608.
138. Shinoda Y, Kozaki K, Imoto I, et al. Association of KLK5 overexpression with invasiveness of urinary bladder carcinoma cells. *Cancer Sci* 2007; 98: 1078-1086.
139. Chen JC, Ho CL, Tsai HW, et al. Immunohistochemical detection of prostate-specific antigen expression in primary urothelial carcinoma of the urinary bladder. *Anticancer Res* 2008; 28: 4149-4154.
140. Kunju LP, Mehra R, Snyder M, et al. Prostate-specific antigen high-molecular-weight cytokeratin (clone 34betaE12) and/or p63: an optimal immunohistochemical panel to distinguish poorly differentiated prostate adenocarcinoma from urothelial carcinoma. *Am J Clin Pathol* 2006; 125: 675-681.
141. Lane Z, Hansel DE, Epstein JI. Immunohistochemical expression of prostatic antigens in adenocarcinoma and villous adenoma of the urinary bladder. *Am J Surg Pathol* 2008; 32: 1322-1326.
142. Clements J, Mukhtar A. Tissue kallikrein and the bradykinin B2 receptor are expressed in endometrial and prostate cancers. *Immunopharmacology* 1997; 36: 217-220.
143. Clements J, Mukhtar A, Ehrlich A, et al. Glandular kallikrein gene expression in the human uterus. *Braz J Med Biol Res* 1994; 27: 1855-1863.
144. Clements J, Mukhtar A. Glandular kallikreins and prostate-specific antigen are expressed in the human endometrium. *J Clin Endocrinol Metab* 1994; 78: 1536-1539.
145. Santin AD, Zhan F, Cané S, et al. Gene expression fingerprint of uterine serous papillary carcinoma: identification of novel molecular markers for uterine serous cancer diagnosis and therapy. *Br J Cancer* 2005; 92: 1561-1573.
146. Seliger E, Kaltwasser P, Röpke F. Determination of prostate-specific antigen (PSA) in cytosol of breast tumours and human endometrium--new diagnostic approaches. *Zentralbl Gynakol* 1998; 120: 172-175.
147. Kucera E, Kainz C, Tempfer C, et al. Prostate specific antigen PSA in breast and ovarian cancer. *Anticancer Res* 1997; 17: 4735-4737.
148. Yu H, Diamandis EP, Levesque M, et al. Expression of the prostate-specific antigen gene by a primary ovarian carcinoma. *Cancer Res* 1995; 55: 1603-1606.
149. Davidson B, Xi Z, Klock TI, et al. Kallikrein 4 expression is up-regulated in epithelial ovarian carcinoma cells in effusions. *Am J Clin Pathol* 2005; 123: 360-368.
150. Hibbs K, Skubitz KM, Pambuccian SE, et al. Differential gene expression in ovarian carcinoma: identification of potential biomarkers. *Am J Pathol* 2004; 165: 397-414.
151. Youssef GM, Scorilas A, Kyriakopoulou LG, et al. Human kallikrein gene 5 KLK5 expression by quantitative PCR: an independent indicator of poor prognosis in breast cancer. *Clin Chem* 2002; 48: 1241-1250.
152. Anisowicz A, Sotiropoulou G, Stenman G, et al. Novel protease homolog differentially expressed in breast and ovarian cancer. *Mol Med* 1996; 2: 624-626.
153. Gyorffy B, Lanczky A, Szallasi Z. Implementing an online tool for genome-wide validation of survival-associated biomarkers in ovarian-cancer using microarray data from 1287 patients. *Endocr Relat Cancer* 2012; 19: 197.
154. Hu CJ, Zhang F, Chen YJ, et al. Correlation of hK6 expression with clinicopathological features and prognosis in epithelial ovarian cancer. *Zhonghua Zhong Liu Za Zhi* 2009; 31: 520-523.
155. Lu KH, Patterson AP, Wang L, et al. Selection of potential markers for epithelial ovarian cancer with gene expression arrays and recursive descent partition analysis. *Clin Cancer Res* 2004; 10: 3291-3300.
156. Ni X, Zhang W, Huang KC, et al. Characterisation of human kallikrein 6/protease M expression in ovarian cancer. *Br J Cancer* 2004; 91: 725-731.
157. Shan SJ, Scorilas A, Katsaros D, et al. Transcriptional upregulation of human tissue kallikrein 6 in ovarian cancer: clinical and mechanistic aspects. *Br J Cancer* 2007; 96: 362-372.
158. Welsh JB, Sapinoso LM, Kern SG, et al. Large-scale delineation of secreted protein biomarkers overexpressed in cancer tissue and serum. *Proc Natl Acad Sci USA* 2003; 100: 3410-3415.
159. Youssef GM, Borgono CA, White NM, et al. In silico analysis of the human kallikrein gene 6. *Tumour Biol* 2004; 25: 282-289.
160. Psyrri A, Kountourakis P, Scorilas A, et al. Human tissue kallikrein 7 a novel biomarker for advanced ovarian carcinoma using a novel in situ quantitative method of protein expression. *Ann Oncol* 2008; 19: 1271-1277.
161. Youssef GM, Fracchioli S, Scorilas A, et al. Steroid hormone regulation and prognostic value of the human kallikrein gene 14 in ovarian cancer. *Am J Clin Pathol* 2003; 119: 346-355.
162. Batra J, Nagle CM, O'Mara T, et al. A kallikrein 15 KLK15 single nucleotide polymorphism located close to a novel exon shows evidence of association with poor ovarian cancer survival. *BMC Cancer* 2011; 11: 119.
163. Dong Y, Matigian N, Harvey TJ, et al. Tissue-specific promoter utilisation of the kallikrein-related peptidase genes KLK5 and KLK7 and cellular localisation of the encoded proteins suggest roles in exocrine pancreatic function. *Biol Chem* 2008; 389: 99-109.
164. Harvey TJ, Hooper JD, Myers SA, et al. Tissue-specific expression patterns and fine mapping of the human kallikrein (KLK) locus on proximal 19q13.4. *J Biol Chem* 2000; 275: 37397-37406.
165. Hooper JD, Bui LT, Rae FK, et al. Identification and characterisation of KLK14 a novel kallikrein serine protease gene located on human chromosome 19q13.4 and expressed in prostate and skeletal muscle. *Genomics* 2001; 73: 117-122.
166. Kishi T, Soosaipillai A, Grass L, et al. Development of an immunofluorometric assay and quantification of human kallikrein 7 in tissue extracts and biological fluids. *Clin Chem* 2004; 50: 709-716.
167. Obiezu CV, Shan SJ, Soosaipillai A, et al. Human kallikrein 4: quantitative study in tissues and evidence for its secretion into biological fluids. *Clin Chem* 2005; 51: 1432-1442.
168. Youssef GM, Chang A, Scorilas A, et al. Genomic organisation of the human kallikrein gene family on chromosome 19q13.3-q13.4. *Biochem Biophys Res Commun* 2000; 276: 125-133.
169. Youssef GM, Obiezu CV, Luo LY, et al. Prostase/KLK-L1 is a new member of the human kallikrein gene family expressed in prostate and breast tissues and is hormonally regulated. *Cancer Res* 1999; 59: 4252-4256.
170. Becker C, Noldus J, Diamandis E, et al. The role of molecular forms of prostate-specific antigen (PSA or hK3) and of human glandular kallikrein 2 (hK2) in the diagnosis and monitoring of prostate cancer and in extra-prostatic disease. *Crit Rev Clin Lab Sci* 2001; 38: 357-399.
171. Kwiatkowski MK, Recker F, Piironen T, et al. In prostatism patients the ratio of human glandular kallikrein to free PSA improves the discrimination between prostate cancer and benign hyperplasia within the diagnostic "gray zone" of total PSA 4 to 10 ng/mL. *Urology* 1998; 52: 360-365.
172. Magklara A, Scorilas A, Catalona WJ, et al. The combination of human glandular kallikrein and free prostate-specific antigen (PSA) enhances discrimination between prostate cancer and benign prostatic hyperplasia in patients with moderately increased total PSA. *Clin Chem* 1999; 45: 1960-1966.
173. Ylikoski A, Pettersson K, Nurmi J, et al. Simultaneous quantification of prostate-specific antigen and human glandular kallikrein 2 mRNA in blood samples from patients with prostate cancer and benign disease. *Clin Chem* 2002; 48: 1265-1271.
174. Stephan C, Jung K, Lein M, et al. PSA and other tissue kallikreins for prostate cancer detection. *Eur J Cancer* 2007; 43: 1918-1926.
175. Steuber T, Vickers A, Haese A, et al. Free PSA isoforms and intact and cleaved forms of urokinase plasminogen activator receptor in serum improve selection of patients for prostate cancer biopsy. *Int J Cancer* 2007; 120: 1499-1504.
176. Steuber T, Vickers AJ, Serio AM, et al. Comparison of free and total forms of serum human kallikrein 2 and prostate-specific antigen for prediction of locally advanced and recurrent prostate cancer. *Clin Chem* 2007; 53: 233-240.
177. Avgeris M, Stravodimos K, Scorilas A. Kallikrein-related peptidase 4 gene (KLK4) in prostate tumours: quantitative expression analysis and evaluation of its clinical significance. *Prostate* 2011; 71: 1780-1789.
178. Dong Y, Bui LT, Odorico DM, et al. Compartmentalized expression of kallikrein 4 (KLK4/hK4) isoforms in prostate cancer: nuclear cytoplasmic and secreted forms. *Endocr Relat Cancer* 2005; 12: 875-889.

- 179Klokk TI, Kilander A, Xi Z, et al. Kallikrein 4 is a proliferative factor that is over-expressed in prostate cancer. *Cancer Res* 2007; 67: 5221-52230.
- 180Petraki CD, Gregorakis AK, Papanastasiou PA, et al. Immunohistochemical localisation of human kallikreins 6 10 and 13 in benign and malignant prostatic tissues. *Prostate Cancer Prostatic Dis* 2003; 6: 223-227.
- 181Petraki CD, Karavana VN, Revelos KI, et al. Immunohistochemical localisation of human kallikreins 6 and 10 in pancreatic islets. *Histochem J* 2002; 34: 313-322.
- 182Ramsay AJ, Dong Y, Hunt ML, et al. Kallikrein-related peptidase 4 (KLK4) initiates intracellular signaling via protease-activated receptors (PARs; KLK4 and PAR-2 are co-expressed during prostate cancer progression. *J Biol Chem* 2008; 283: 12293-12304.
- 183Seiz L, Kotzsch M, Grebentchikov NI, et al. Polyclonal antibodies against kallikrein-related peptidase 4 (KLK4): immunohistochemical assessment of KLK4 expression in healthy tissues and prostate cancer. *Biol Chem* 2010; 391: 391-401.
- 184Stephenson SA, Verity K, Ashworth LK, et al. Localisation of a new prostate-specific antigen-related serine protease gene KLK4 is evidence for an expanded human kallikrein gene family cluster on chromosome 19q13.3-13.4. *J Biol Chem* 1999; 274: 23210-23214.
- 185Korbakis D, Gregorakis AK, Scorilas A. Quantitative analysis of human kallikrein 5 (KLK5) expression in prostate needle biopsies: an independent cancer biomarker. *Clin Chem* 2009; 55: 904-913.
- 186Yousef GM, Scorilas A, Chang A, et al. Down-regulation of the human kallikrein gene 5 (KLK5) in prostate cancer tissues. *Prostate* 2002; 51: 126-132.
- 187Jamaspishvili T, Scorilas A, Kral M, et al. Immunohistochemical localisation and analysis of kallikrein-related peptidase 7 and 11 expression in paired cancer and benign foci in prostate cancer patients. *Neoplasia* 2011; 58: 298-303.
- 188Xuan Q, Yang X, Mo L, et al. Expression of the serine protease kallikrein 7 and its inhibitor antileukoprotease is decreased in prostate cancer. *Arch Pathol Lab Med* 2008; 132: 1796-1801.
- 189Yousef GM, Scorilas A, Magklara A, et al. The KLK7 (PRSS6) gene encoding for the stratum corneum chymotryptic enzyme is a new member of the human kallikrein gene family - genomic characterisation mapping tissue expression and hormonal regulation. *Gene* 2000; 254: 119-128.
- 190Kishi T, Grass L, Soosaipillai A, et al. Human kallikrein 8: immunoassay development and identification in tissue extracts and biological fluids. *Clin Chem* 2003; 49: 87-96.
- 191Bi X, He H, Ye Y, et al. Association of TMPRSS2 and KLK11 gene expression levels with clinical progression of human prostate cancer. *Med Oncol* 2010; 27: 145-151.
- 192Diamandis EP, Okui A, Mitsui S, et al. Human kallikrein 11: a new biomarker of prostate and ovarian carcinoma. *Cancer Res* 2002; 62: 295-300.
- 193Luo LY, Shan SJ, Elliott MB, et al. Purification and characterisation of human kallikrein 11 a candidate prostate and ovarian cancer biomarker from seminal plasma. *Clin Cancer Res* 2006; 12: 742-750.
- 194Scorilas A, Gregorakis AK. mRNA expression analysis of human kallikrein 11 (KLK11) may be useful in the discrimination of benign prostatic hyperplasia from prostate cancer after needle prostate biopsy. *Biol Chem* 2006; 387: 789-793.
- 195Stavropoulou P, Gregorakis AK, Plebani M, et al. Expression analysis and prognostic significance of human kallikrein 11 in prostate cancer. *Clin Chim Acta* 2005; 357: 190-195.
- 196Memari N, Diamandis EP, Earle T, et al. Human kallikrein-related peptidase 12: antibody generation and immunohistochemical localisation in prostatic tissues. *Prostate* 2007; 67: 1465-1474.
- 197Nakamura T, Scorilas A, Stephan C, et al. The usefulness of serum human kallikrein 11 for discriminating between prostate cancer and benign prostatic hyperplasia. *Cancer Res* 2003; 63: 6543-6546.
- 198Nakamura T, Stephan C, Scorilas A, et al. Quantitative analysis of hippostasin/KLK11 gene expression in cancerous and noncancerous prostatic tissues. *Urology* 2003; 61: 1042-1046.
- 199Rabien A, Fritzsche F, Jung M, et al. High expression of KLK14 in prostatic adenocarcinoma is associated with elevated risk of prostate-specific antigen relapse. *Tumour Biol* 2008; 29: 1-8.
- 200Yousef GM, Stephan C, Scorilas A, et al. Differential expression of the human kallikrein gene 14 (KLK14) in normal and cancerous prostatic tissues. *Prostate* 2003; 56: 287-292.
- 201Cumming AD, Walsh T, Wojtacha D, et al. Expression of tissue kallikrein in human kidney. *Clin Sci (Lond)* 1994; 87: 5-11.
- 202Rae F, Bulmer B, Nicol D, et al. The human tissue kallikreins (KLKs 1-3) and a novel KLK1 mRNA transcript are expressed in a renal cell carcinoma cDNA library. *Immunopharmacology* 1999; 45: 83-88.
- 203Kishi T, Cloutier SM, Kundig C, et al. Activation and enzymatic characterisation of recombinant human kallikrein 8. *Biol Chem* 2006; 387: 723-731.
- 204Kontos CK, Scorilas A. Kallikrein-related peptidases (KLKs): a gene family of novel cancer biomarkers. *Clin Chem Lab Med* 2012; 50: 1877-1891.
- 205Sotiropoulou G, Pampalakis G. Targeting the kallikrein-related peptidases for drug development. *Trends Pharmacol Sci* 2012; 33: 623-634.
- 206Beaufort N, Plaza K, Utzschneider D, Schwarz A, Burkhart JM, Creutzburg S, Debela M, Schmitt M, Ries C, Magdolen V. Interdependence of kallikrein-related peptidases in proteolytic networks. *Biol Chem* 2010; 391: 581-587.
- 207Debela M, Beaufort N, Magdolen V, Schechter NM, Craik CS, Schmitt M, Bode W, Goettig P. Structures and specificity of the human kallikrein-related peptidases KLK 4, 5, 6, and 7. *Biol Chem*; 389: 623-632.